

CORRECTION TO A THEOREM OF SCHOENBERG

CARL JOHAN RAGNARSSON, WESLEY WAI SUEN, AND DAVID G. WAGNER

ABSTRACT. A well-known theorem of Schoenberg states that if $f(z)$ generates a PF_r sequence then $1/f(-z)$ generates a PF_r sequence. We give two counterexamples which show that this is not true, and give a correct version of the theorem. In the infinite limit the result is sound: if $f(z)$ generates a PF sequence then $1/f(-z)$ generates a PF sequence.

1. THE BAD NEWS.

Theorem 1.2 in Chapter 8 of Karlin's book [2] implies the following:

Theorem A. *Let $f(z) = 1 + \sum_{n=1}^{\infty} a_n z^n$ and $g(z) = 1 + \sum_{n=1}^{\infty} b_n z^n$ be power series with real coefficients such that $g(z) = 1/f(-z)$. For any positive integer r , the Toeplitz matrix of f is totally positive up to order r if and only if the Toeplitz matrix of g is totally positive up to order r .*

Theorem A first appears in work of Schoenberg *et al.* in the early 1950s [1, 4, 5]. The bad news is that Theorem A is false.

First let's review the definitions. For a power series $f(z) = \sum_{n=0}^{\infty} a_n z^n$, the *Toeplitz matrix of f* is the infinite matrix $T[f]$, indexed by pairs of integers, with entries

$$T[f]_{ij} := \begin{cases} a_{j-i} & \text{if } j - i \geq 0, \\ 0 & \text{if } j - i < 0. \end{cases}$$

An infinite matrix M is *totally positive up to order r* when every minor of M of order at most r is nonnegative. This condition is abbreviated TP_r . If M is TP_r for all r then M is *totally positive*, abbreviated TP.

The matrix $T[f]$ is TP_1 if and only if the coefficients of $f(z)$ are nonnegative. If $T[f]$ is TP_2 then the sequence of coefficients $a_0, a_1, \dots$ has *no internal zeros*: if $0 \leq h < i < j$ and $a_h a_j \neq 0$, then $a_i \neq 0$.

1991 *Mathematics Subject Classification.* 15A48; 15A45, 15A57, 05E05.

Key words and phrases. Total positivity, Pólya frequency sequence, skew Schur function.

0. Also, if $T[f]$ is TP_2 then the sequence of coefficients $a_0, a_1, \dots$ is *logarithmically concave*: if $j \geq 1$ then $a_j^2 \geq a_{j-1}a_{j+1}$. Nonnegativity of the remaining 2-by-2 minors of $T[f]$ follow from these two conditions. That is, the Toeplitz matrix $T[f]$ is TP_2 if and only if the sequence of coefficients $a_0, a_1, \dots$ is nonnegative, has no internal zeros, and is logarithmically concave.

Our first counterexample to Theorem A is the polynomial $f(z) = 1 + 4z + 3z^2 + z^3$. By the preceding paragraph, one sees easily that $T[f]$ is TP_2 . Elementary calculation with linear recurrence relations yields

$$g(z) = \frac{1}{1 - 4z + 3z^2 - z^3} = 1 + 4z + 13z^2 + 41z^3 + 129z^4 + 406z^5 + \dots$$

Since $129^2 - 41 \cdot 406 = -5 < 0$, the Toeplitz matrix $T[g]$ is evidently not TP_2 . Theorem A is false.

With hindsight, one notices that the coefficients of $f(z) = 1 + z + 2z^2$ are nonnegative, but that

$$g(z) = \frac{1}{1 - z + 2z^2} = 1 + z - z^2 - 3z^3 - z^4 + 5z^5 + \dots$$

has negative coefficients. Thus, $T[f]$ is TP_1 while $T[g]$ is not TP_1 . This is a rather glaring counterexample to Theorem A.

2. THE GOOD NEWS.

The good news is that Theorem A can be fixed.

To do this we need a few facts about symmetric functions – see Macdonald [3] for details. Let $e_1, e_2, \dots$ and $h_1, h_2, \dots$ be indeterminates which are algebraically independent over the field $\mathbb{Q}$ of rational numbers, and form the generic power series $E(t) := 1 + \sum_{n=1}^{\infty} e_n t^n$ and $H(t) := 1 + \sum_{n=1}^{\infty} h_n t^n$. By imposing the single relation $E(t) = H(-t)^{-1}$ one can determine each e_n as a polynomial in the h_n -s, and conversely. The indeterminates $\{h_n\}$ remain algebraically independent over $\mathbb{Q}$, as do the indeterminates $\{e_n\}$. The ring Λ of polynomials with integer coefficients in these indeterminates is the *ring of symmetric functions*.

Since the indeterminates $\{h_n\}$ are algebraically independent and generate Λ , a homomorphism $\varphi : \Lambda \rightarrow R$ from Λ to another ring R is determined by its values $\{\varphi(h_n)\}$. A real power series $f(z) = 1 + \sum_{n=0}^{\infty} a_n z^n$ determines such a homomorphism $\varphi_f : \Lambda \rightarrow \mathbb{R}$ by $\varphi_f(h_n) := a_n$. Notice that if $g(z) = 1 + \sum_{n=1}^{\infty} b_n z^n$ is such that $g(z) = 1/f(-z)$ then $\varphi_f(e_n) = b_n$ and $\varphi_g(e_n) = a_n$.

The set of all integer partitions, partially ordered by inclusion of Ferrers diagrams, is called *Young's lattice* and denoted by $\mathcal{Y}$. For $\mu \leq \lambda$

in $\mathcal{Y}$ there is a symmetric function $s_{\lambda/\mu}$ called a *skew Schur function*. Without loss of generality we may assume not only that $\mu \leq \lambda$ in $\mathcal{Y}$, but also that μ has strictly fewer parts than λ and that the largest part of μ is strictly smaller than the largest part of λ . We will denote this relation by $\mu \prec \lambda$ in $\mathcal{Y}$. The formulae we need are the Jacobi–Trudy formula and its dual form:

$$s_{\lambda'/\mu'} = \det(e_{\lambda_i - i + j - \mu_j}) \quad \text{and} \quad s_{\lambda/\mu} = \det(h_{\lambda_i - i + j - \mu_j}).$$

The order of these determinants is the number of parts of λ . The notation λ' denotes the partition conjugate to λ . If $f(z)$ and $g(z)$ are real power series such that $g(z) = 1/f(-z)$ then

$$\varphi_f(s_{\lambda/\mu}) = \varphi_g(s_{\lambda'/\mu'}) \quad \text{and} \quad \varphi_g(s_{\lambda/\mu}) = \varphi_f(s_{\lambda'/\mu'}).$$

Consider the submatrix M of $T[f]$ supported on rows $\{i_1 < i_2 < \cdots < i_r\}$ and columns $\{j_1 < j_2 < \cdots < j_r\}$. If $j_k < i_k$ for any $1 \leq k \leq r$ then $\det(M) = 0$, so we may assume that $j_k \geq i_k$ for all $1 \leq k \leq r$. If $j_1 = i_1$ or $j_r = i_r$ then $\det(M)$ reduces by Laplace expansion to a smaller minor of $T[f]$. Thus we may assume as well that $j_1 > i_1$ and $j_r > i_r$. A minor satisfying all these conditions is called an *essential minor* of $T[f]$. It is clear that $T[f]$ is TP_r if and only if every essential minor of $T[f]$ of order at most r is nonnegative.

Every essential minor of $T[f]$ has the form $\varphi_f(s_{\lambda/\mu}) = \det(a_{\lambda_i - i + j - \mu_j})$ for some $\mu \prec \lambda$ in $\mathcal{Y}$. To see this, let $\det(M)$ be an essential minor of $T[f]$ supported on rows $\{i_1 < i_2 < \cdots < i_r\}$ and columns $\{j_1 < j_2 < \cdots < j_r\}$. For each $1 \leq k \leq r$ let $\lambda_k := j_r - i_k + k - r$. The inequalities $\lambda_1 \geq \cdots \lambda_r > 0$ are easily seen, so that λ is an integer partition with r parts. For each $1 \leq k \leq r$ let $\mu_k := j_r - j_k + k - r$. One can check that μ is an integer partition with at most $r - 1$ parts, that $\mu \prec \lambda$ in $\mathcal{Y}$, and that $\det(M) = \det(a_{\lambda_i - i + j - \mu_j})$. This construction can be reversed, so that every $\varphi_f(s_{\lambda/\mu})$ is an essential minor of $T[f]$. In this way the skew Schur functions can be regarded as “generic essential Toeplitz minors”.

The order of the minor $\varphi_f(s_{\lambda/\mu})$ of $T[f]$ is the number of parts of λ . This implies the following:

(a) The Toeplitz matrix $T[f]$ is TP_r if and only if $\varphi_f(s_{\lambda/\mu}) \geq 0$ for all $\mu \prec \lambda$ in $\mathcal{Y}$ for which λ has at most r parts.

Similarly,

(b) The Toeplitz matrix $T[g]$ is TP_r if and only if $\varphi_g(s_{\lambda/\mu}) \geq 0$ for all $\mu \prec \lambda$ in $\mathcal{Y}$ for which λ has at most r parts.

If $g(z) = 1/f(-z)$ then, since $\varphi_g(s_{\lambda/\mu}) = \varphi_f(s_{\lambda'/\mu'})$, condition (b) is equivalent to

(c) The Toeplitz matrix $T[g]$ is TP_r if and only if $\varphi_f(s_{\lambda'/\mu'}) \geq 0$ for all $\mu \prec \lambda$ in $\mathcal{Y}$ for which λ has at most r parts.

Or, in other words,

(d) The Toeplitz matrix $T[g]$ is TP_r if and only if $\varphi_f(s_{\lambda/\mu}) \geq 0$ for all $\mu \prec \lambda$ in $\mathcal{Y}$ for which λ has largest part at most r .

Comparing (a) and (d) we see that the two conditions in Theorem A are closely related, but not equivalent.

Interpreting $\varphi_f(s_{\lambda/\mu})$ as a minor of $T[f]$, bounding the number of parts of λ corresponds to bounding the order of the minor. What corresponds to bounding the largest part of λ ? For the submatrix M of $T[f]$ supported on rows $\{i_1 < i_2 < \cdots < i_r\}$ and columns $\{j_1 < j_2 < \cdots < j_r\}$, define the *level* of M to be $\ell := j_r - i_1 + 1 - r$. The level of a minor of $T[f]$ is the level of the submatrix of which it is the determinant. The Toeplitz matrix $T[f]$ is *totally positive up to level* ℓ when every minor of $T[f]$ of level at most ℓ is nonnegative. This condition is abbreviated TP'_ℓ . If $T[f]$ is TP'_ℓ for all ℓ then $T[f]$ is totally positive, TP .

Theorem B. *Let $f(z) = 1 + \sum_{n=1}^{\infty} a_n z^n$ and $g(z) = 1 + \sum_{n=1}^{\infty} b_n z^n$ be power series with real coefficients such that $g(z) = 1/f(-z)$. For any positive integer r , $T[f]$ is totally positive up to level r if and only if $T[g]$ is totally positive up to order r .*

Notice that in the limit as $r \rightarrow \infty$ we get the equivalence: $T[f]$ is TP if and only if $T[g]$ is TP . This is the most important consequence of Theorem A in the literature, and it is a huge relief that it survives.

REFERENCES

- [1] M. Aissen, I.J. Schoenberg, and A. Whitney, *On the generating functions of totally positive sequences*, Proc. Nat. Acad. Sci. U.S.A. **37** (1952), 303–307.
- [2] S. Karlin, “Total Positivity, vol. I,” Stanford U.P., Stanford CA, 1968.
- [3] I.G. Macdonald, “Symmetric Functions and Hall Polynomials” (2nd ed.), Oxford U.P., Oxford, 1995.
- [4] I.J. Schoenberg, *On smoothing operations and their generating functions*, Bull. Amer. Math. Soc. **59** (1953), 199–230.
- [5] I.J. Schoenberg, *On the zeros of the generating functions of multiply positive sequences and functions*, Ann. Math. **62** (1955), 447–471.

DEPARTMENT OF COMBINATORICS AND OPTIMIZATION, UNIVERSITY OF WATERLOO, WATERLOO, ONTARIO, CANADA N2L 3G1

E-mail address: wwsuen@math.uwaterloo.ca

E-mail address: dgwagner@math.uwaterloo.ca